Lattices and Orders in Isabelle/HOL

Markus Wenzel
TU München

October 24, 2026

Abstract

We consider abstract structures of orders and lattices. Many fundamental concepts of lattice theory are developed, including dual structures, properties of bounds versus algebraic laws, lattice operations versus set-theoretic ones etc. We also give example instantiations of lattices and orders, such as direct products and function spaces. Well-known properties are demonstrated, like the Knaster-Tarski Theorem for complete lattices.

This formal theory development may serve as an example of applying Isabelle/HOL to the domain of mathematical reasoning about "axiomatic" structures. Apart from the simply-typed classical set-theory of HOL, we employ Isabelle's system of axiomatic type classes for expressing structures and functors in a light-weight manner. Proofs are expressed in the Isar language for readable formal proof, while aiming at its "best-style" of representing formal reasoning.

Contents

1 Orders **3**
1.1 Ordered structures . . . 3
1.2 Duality . . . 3
1.3 Transforming orders . . . 5
1.3.1 Duals . . . 5
1.3.2 Binary products . . . 6
1.3.3 General products . . . 7

2 Bounds **8**
2.1 Infimum and supremum . . . 8
2.2 Duality . . . 10
2.3 Uniqueness . . . 10
2.4 Related elements . . . 11
2.5 General versus binary bounds . . . 12
2.6 Connecting general bounds . . . 13

3 Lattices **14**
3.1 Lattice operations 14
3.2 Duality 16
3.3 Algebraic properties 17
3.4 Order versus algebraic structure 19
3.5 Example instances 20
3.5.1 Linear orders 20
3.5.2 Binary products 21
3.5.3 General products 23
3.6 Monotonicity and semi-morphisms 24

4 Complete lattices **26**
4.1 Complete lattice operations 26
4.2 The Knaster-Tarski Theorem 27
4.3 Bottom and top elements 29
4.4 Duality 30
4.5 Complete lattices are lattices 31
4.6 Complete lattices and set-theory operations 32

1 Orders

theory *Orders* **imports** *Main* **begin**

1.1 Ordered structures

We define several classes of ordered structures over some type $'a$ with relation $\sqsubseteq :: 'a \Rightarrow 'a \Rightarrow bool$. For a *quasi-order* that relation is required to be reflexive and transitive, for a *partial order* it also has to be anti-symmetric, while for a *linear order* all elements are required to be related (in either direction).

class *leq* =
  fixes *leq* :: $'a \Rightarrow 'a \Rightarrow bool$ (**infixl** ‹$\sqsubseteq$› *50*)

class *quasi-order* = *leq* +
  assumes *leq-refl* [*intro?*]: $x \sqsubseteq x$
  assumes *leq-trans* [*trans*]: $x \sqsubseteq y \Longrightarrow y \sqsubseteq z \Longrightarrow x \sqsubseteq z$

class *partial-order* = *quasi-order* +
  assumes *leq-antisym* [*trans*]: $x \sqsubseteq y \Longrightarrow y \sqsubseteq x \Longrightarrow x = y$

class *linear-order* = *partial-order* +
  assumes *leq-linear*: $x \sqsubseteq y \vee y \sqsubseteq x$

lemma *linear-order-cases*:
    $((x::'a::linear\text{-}order) \sqsubseteq y \Longrightarrow C) \Longrightarrow (y \sqsubseteq x \Longrightarrow C) \Longrightarrow C$
  by (*insert leq-linear*) *blast*

1.2 Duality

The *dual* of an ordered structure is an isomorphic copy of the underlying type, with the $\sqsubseteq$ relation defined as the inverse of the original one.

datatype $'a$ *dual* = *dual* $'a$

primrec *undual* :: $'a$ *dual* $\Rightarrow$ $'a$ **where**
  undual-dual: *undual* (*dual* x) = x

instantiation *dual* :: (*leq*) *leq*
begin

definition
  leq-dual-def: $x' \sqsubseteq y' \equiv$ *undual* $y' \sqsubseteq$ *undual* x'

instance ..

end

lemma *undual-leq* [*iff?*]: (*undual* $x' \sqsubseteq$ *undual* y') = ($y' \sqsubseteq x'$)
  by (*simp add*: *leq-dual-def*)

lemma *dual-leq* [*iff?*]: $(dual\ x \sqsubseteq dual\ y) = (y \sqsubseteq x)$
by (*simp add*: *leq-dual-def*)

Functions *dual* and *undual* are inverse to each other; this entails the following fundamental properties.

lemma *dual-undual* [*simp*]: $dual\ (undual\ x') = x'$
by (*cases* x') *simp*

lemma *undual-dual-id* [*simp*]: $undual\ o\ dual = id$
by (*rule ext*) *simp*

lemma *dual-undual-id* [*simp*]: $dual\ o\ undual = id$
by (*rule ext*) *simp*

Since *dual* (and *undual*) are both injective and surjective, the basic logical connectives (equality, quantification etc.) are transferred as follows.

lemma *undual-equality* [*iff?*]: $(undual\ x' = undual\ y') = (x' = y')$
by (*cases* x', *cases* y') *simp*

lemma *dual-equality* [*iff?*]: $(dual\ x = dual\ y) = (x = y)$
by *simp*

lemma *dual-ball* [*iff?*]: $(\forall x \in A.\ P\ (dual\ x)) = (\forall x' \in dual\ `\ A.\ P\ x')$
proof
assume *a*: $\forall x \in A.\ P\ (dual\ x)$
show $\forall x' \in dual\ `\ A.\ P\ x'$
proof
fix x' **assume** x': $x' \in dual\ `\ A$
have $undual\ x' \in A$
proof −
from x' **have** $undual\ x' \in undual\ `\ dual\ `\ A$ **by** *simp*
thus $undual\ x' \in A$ **by** (*simp add*: *image-comp*)
qed
with *a* **have** $P\ (dual\ (undual\ x'))$ **..**
also have $\ldots = x'$ **by** *simp*
finally show $P\ x'$ **.**
qed
next
assume *a*: $\forall x' \in dual\ `\ A.\ P\ x'$
show $\forall x \in A.\ P\ (dual\ x)$
proof
fix x **assume** $x \in A$
hence $dual\ x \in dual\ `\ A$ **by** *simp*
with *a* **show** $P\ (dual\ x)$ **..**
qed
qed

lemma *range-dual* [*simp*]: *surj dual*
proof −
  have $\bigwedge x'$. *dual* (*undual* x') = x' **by** *simp*
  thus *surj dual* **by** (*rule surjI*)
qed

lemma *dual-all* [*iff?*]: $(\forall\, x.\ P\ (\mathit{dual}\ x)) = (\forall\, x'.\ P\ x')$
proof −
  have $(\forall\, x \in \mathit{UNIV}.\ P\ (\mathit{dual}\ x)) = (\forall\, x' \in \mathit{dual}\ `\ \mathit{UNIV}.\ P\ x')$
    by (*rule dual-ball*)
  thus *?thesis* **by** *simp*
qed

lemma *dual-ex*: $(\exists\, x.\ P\ (\mathit{dual}\ x)) = (\exists\, x'.\ P\ x')$
proof −
  have $(\forall\, x.\ \neg\ P\ (\mathit{dual}\ x)) = (\forall\, x'.\ \neg\ P\ x')$
    by (*rule dual-all*)
  thus *?thesis* **by** *blast*
qed

lemma *dual-Collect*: $\{\mathit{dual}\ x |\ x.\ P\ (\mathit{dual}\ x)\} = \{x'.\ P\ x'\}$
proof −
  have $\{\mathit{dual}\ x |\ x.\ P\ (\mathit{dual}\ x)\} = \{x'.\ \exists\, x''.\ x' = x'' \wedge P\ x''\}$
    by (*simp only*: *dual-ex* [*symmetric*])
  thus *?thesis* **by** *blast*
qed

1.3 Transforming orders

1.3.1 Duals

The classes of quasi, partial, and linear orders are all closed under formation of dual structures.

instance *dual* :: (*quasi-order*) *quasi-order*
proof
  fix x' y' z' :: *'a*::*quasi-order dual*
  have *undual* $x' \sqsubseteq$ *undual* x' **.. thus** $x' \sqsubseteq x'$ **..**
  assume $y' \sqsubseteq z'$ **hence** *undual* $z' \sqsubseteq$ *undual* y' **..**
  also assume $x' \sqsubseteq y'$ **hence** *undual* $y' \sqsubseteq$ *undual* x' **..**
  finally show $x' \sqsubseteq z'$ **..**
qed

instance *dual* :: (*partial-order*) *partial-order*
proof
  fix x' y' :: *'a*::*partial-order dual*
  assume $y' \sqsubseteq x'$ **hence** *undual* $x' \sqsubseteq$ *undual* y' **..**
  also assume $x' \sqsubseteq y'$ **hence** *undual* $y' \sqsubseteq$ *undual* x' **..**
  finally show $x' = y'$ **..**
qed

instance *dual* :: (*linear-order*) *linear-order*
proof
  fix x' y' :: $'a$::*linear-order dual*
  show $x' \sqsubseteq y' \vee y' \sqsubseteq x'$
  proof (*rule linear-order-cases*)
    assume *undual* $y' \sqsubseteq$ *undual* x'
    hence $x' \sqsubseteq y'$ **.. thus** *?thesis* **..**
  next
    assume *undual* $x' \sqsubseteq$ *undual* y'
    hence $y' \sqsubseteq x'$ **.. thus** *?thesis* **..**
  qed
qed

1.3.2 Binary products

The classes of quasi and partial orders are closed under binary products. Note that the direct product of linear orders need *not* be linear in general.

instantiation *prod* :: (*leq*, *leq*) *leq*
begin

definition
  leq-prod-def: $p \sqsubseteq q \equiv$ *fst* $p \sqsubseteq$ *fst* $q \wedge$ *snd* $p \sqsubseteq$ *snd* q

instance ..

end

lemma *leq-prodI* [*intro?*]:
    fst $p \sqsubseteq$ *fst* $q \Longrightarrow$ *snd* $p \sqsubseteq$ *snd* $q \Longrightarrow p \sqsubseteq q$
  by (*unfold leq-prod-def*) *blast*

lemma *leq-prodE* [*elim?*]:
    $p \sqsubseteq q \Longrightarrow$ (*fst* $p \sqsubseteq$ *fst* $q \Longrightarrow$ *snd* $p \sqsubseteq$ *snd* $q \Longrightarrow C) \Longrightarrow C$
  by (*unfold leq-prod-def*) *blast*

instance *prod* :: (*quasi-order*, *quasi-order*) *quasi-order*
proof
  fix p q r :: $'a$::*quasi-order* $\times$ $'b$::*quasi-order*
  show $p \sqsubseteq p$
  proof
    show *fst* $p \sqsubseteq$ *fst* p **..**
    show *snd* $p \sqsubseteq$ *snd* p **..**
  qed
  assume *pq*: $p \sqsubseteq q$ **and** *qr*: $q \sqsubseteq r$
  show $p \sqsubseteq r$
  proof
    from *pq* **have** *fst* $p \sqsubseteq$ *fst* q **..**
    also from *qr* **have** $\ldots \sqsubseteq$ *fst* r **..**

```
    finally show fst p ⊑ fst r .
    from pq have snd p ⊑ snd q ..
    also from qr have ... ⊑ snd r ..
    finally show snd p ⊑ snd r .
  qed
qed

instance prod :: (partial-order, partial-order) partial-order
proof
  fix p q :: 'a::partial-order × 'b::partial-order
  assume pq: p ⊑ q and qp: q ⊑ p
  show p = q
  proof
    from pq have fst p ⊑ fst q ..
    also from qp have ... ⊑ fst p ..
    finally show fst p = fst q .
    from pq have snd p ⊑ snd q ..
    also from qp have ... ⊑ snd p ..
    finally show snd p = snd q .
  qed
qed
```

1.3.3 General products

The classes of quasi and partial orders are closed under general products (function spaces). Note that the direct product of linear orders need *not* be linear in general.

```
instantiation fun :: (type, leq) leq
begin

definition
  leq-fun-def: f ⊑ g ≡ ∀ x. f x ⊑ g x

instance ..

end

lemma leq-funI [intro?]: (⋀x. f x ⊑ g x) ⟹ f ⊑ g
  by (unfold leq-fun-def) blast

lemma leq-funD [dest?]: f ⊑ g ⟹ f x ⊑ g x
  by (unfold leq-fun-def) blast

instance fun :: (type, quasi-order) quasi-order
proof
  fix f g h :: 'a ⇒ 'b::quasi-order
  show f ⊑ f
  proof
```

fix x **show** $f\,x \sqsubseteq f\,x$ **..**
qed
assume *fg*: $f \sqsubseteq g$ **and** *gh*: $g \sqsubseteq h$
show $f \sqsubseteq h$
proof
fix x **from** *fg* **have** $f\,x \sqsubseteq g\,x$ **..**
also from *gh* **have** $\ldots \sqsubseteq h\,x$ **..**
finally show $f\,x \sqsubseteq h\,x$ **.**
qed
qed

instance *fun* :: (*type*, *partial-order*) *partial-order*
proof
fix $f\,g$ :: $'a \Rightarrow 'b$::*partial-order*
assume *fg*: $f \sqsubseteq g$ **and** *gf*: $g \sqsubseteq f$
show $f = g$
proof
fix x **from** *fg* **have** $f\,x \sqsubseteq g\,x$ **..**
also from *gf* **have** $\ldots \sqsubseteq f\,x$ **..**
finally show $f\,x = g\,x$ **.**
qed
qed

end

2 Bounds

theory *Bounds* **imports** *Orders* **begin**

hide-const (**open**) *inf sup*

2.1 Infimum and supremum

Given a partial order, we define infimum (greatest lower bound) and supremum (least upper bound) wrt. $\sqsubseteq$ for two and for any number of elements.

definition
is-inf :: $'a$::*partial-order* $\Rightarrow$ $'a \Rightarrow 'a \Rightarrow$ *bool* **where**
is-inf $x\ y\ inf = (inf \sqsubseteq x \wedge inf \sqsubseteq y \wedge (\forall z.\ z \sqsubseteq x \wedge z \sqsubseteq y \longrightarrow z \sqsubseteq inf))$

definition
is-sup :: $'a$::*partial-order* $\Rightarrow$ $'a \Rightarrow 'a \Rightarrow$ *bool* **where**
is-sup $x\ y\ sup = (x \sqsubseteq sup \wedge y \sqsubseteq sup \wedge (\forall z.\ x \sqsubseteq z \wedge y \sqsubseteq z \longrightarrow sup \sqsubseteq z))$

definition
is-Inf :: $'a$::*partial-order set* $\Rightarrow$ $'a \Rightarrow$ *bool* **where**
is-Inf $A\ inf = ((\forall x \in A.\ inf \sqsubseteq x) \wedge (\forall z.\ (\forall x \in A.\ z \sqsubseteq x) \longrightarrow z \sqsubseteq inf))$

definition

is-Sup :: *'a::partial-order set* ⇒ *'a* ⇒ *bool* **where**
is-Sup A sup = ((∀ *x* ∈ *A*. *x* ⊑ *sup*) ∧ (∀ *z*. (∀ *x* ∈ *A*. *x* ⊑ *z*) ⟶ *sup* ⊑ *z*))

These definitions entail the following basic properties of boundary elements.

lemma *is-infI* [*intro?*]: *inf* ⊑ *x* ⟹ *inf* ⊑ *y* ⟹
(⋀*z*. *z* ⊑ *x* ⟹ *z* ⊑ *y* ⟹ *z* ⊑ *inf*) ⟹ *is-inf x y inf*
by (*unfold is-inf-def*) *blast*

lemma *is-inf-greatest* [*elim?*]:
is-inf x y inf ⟹ *z* ⊑ *x* ⟹ *z* ⊑ *y* ⟹ *z* ⊑ *inf*
by (*unfold is-inf-def*) *blast*

lemma *is-inf-lower* [*elim?*]:
is-inf x y inf ⟹ (*inf* ⊑ *x* ⟹ *inf* ⊑ *y* ⟹ *C*) ⟹ *C*
by (*unfold is-inf-def*) *blast*

lemma *is-supI* [*intro?*]: *x* ⊑ *sup* ⟹ *y* ⊑ *sup* ⟹
(⋀*z*. *x* ⊑ *z* ⟹ *y* ⊑ *z* ⟹ *sup* ⊑ *z*) ⟹ *is-sup x y sup*
by (*unfold is-sup-def*) *blast*

lemma *is-sup-least* [*elim?*]:
is-sup x y sup ⟹ *x* ⊑ *z* ⟹ *y* ⊑ *z* ⟹ *sup* ⊑ *z*
by (*unfold is-sup-def*) *blast*

lemma *is-sup-upper* [*elim?*]:
is-sup x y sup ⟹ (*x* ⊑ *sup* ⟹ *y* ⊑ *sup* ⟹ *C*) ⟹ *C*
by (*unfold is-sup-def*) *blast*

lemma *is-InfI* [*intro?*]: (⋀*x*. *x* ∈ *A* ⟹ *inf* ⊑ *x*) ⟹
(⋀*z*. (∀ *x* ∈ *A*. *z* ⊑ *x*) ⟹ *z* ⊑ *inf*) ⟹ *is-Inf A inf*
by (*unfold is-Inf-def*) *blast*

lemma *is-Inf-greatest* [*elim?*]:
is-Inf A inf ⟹ (⋀*x*. *x* ∈ *A* ⟹ *z* ⊑ *x*) ⟹ *z* ⊑ *inf*
by (*unfold is-Inf-def*) *blast*

lemma *is-Inf-lower* [*dest?*]:
is-Inf A inf ⟹ *x* ∈ *A* ⟹ *inf* ⊑ *x*
by (*unfold is-Inf-def*) *blast*

lemma *is-SupI* [*intro?*]: (⋀*x*. *x* ∈ *A* ⟹ *x* ⊑ *sup*) ⟹
(⋀*z*. (∀ *x* ∈ *A*. *x* ⊑ *z*) ⟹ *sup* ⊑ *z*) ⟹ *is-Sup A sup*
by (*unfold is-Sup-def*) *blast*

lemma *is-Sup-least* [*elim?*]:
is-Sup A sup ⟹ (⋀*x*. *x* ∈ *A* ⟹ *x* ⊑ *z*) ⟹ *sup* ⊑ *z*

by (*unfold is-Sup-def*) *blast*

lemma *is-Sup-upper* [*dest?*]:
is-Sup A sup $\Longrightarrow$ $x \in A \Longrightarrow x \sqsubseteq$ *sup*
by (*unfold is-Sup-def*) *blast*

2.2 Duality

Infimum and supremum are dual to each other.

theorem *dual-inf* [*iff?*]:
is-inf (*dual x*) (*dual y*) (*dual sup*) = *is-sup x y sup*
by (*simp add*: *is-inf-def is-sup-def dual-all* [*symmetric*] *dual-leq*)

theorem *dual-sup* [*iff?*]:
is-sup (*dual x*) (*dual y*) (*dual inf*) = *is-inf x y inf*
by (*simp add*: *is-inf-def is-sup-def dual-all* [*symmetric*] *dual-leq*)

theorem *dual-Inf* [*iff?*]:
is-Inf (*dual ' A*) (*dual sup*) = *is-Sup A sup*
by (*simp add*: *is-Inf-def is-Sup-def dual-all* [*symmetric*] *dual-leq*)

theorem *dual-Sup* [*iff?*]:
is-Sup (*dual ' A*) (*dual inf*) = *is-Inf A inf*
by (*simp add*: *is-Inf-def is-Sup-def dual-all* [*symmetric*] *dual-leq*)

2.3 Uniqueness

Infima and suprema on partial orders are unique; this is mainly due to antisymmetry of the underlying relation.

theorem *is-inf-uniq*: *is-inf x y inf* $\Longrightarrow$ *is-inf x y inf'* $\Longrightarrow$ *inf* = *inf'*
proof −
assume *inf*: *is-inf x y inf*
assume *inf'*: *is-inf x y inf'*
show *?thesis*
proof (*rule leq-antisym*)
from *inf'* **show** *inf* $\sqsubseteq$ *inf'*
proof (*rule is-inf-greatest*)
from *inf* **show** *inf* $\sqsubseteq x$ **..**
from *inf* **show** *inf* $\sqsubseteq y$ **..**
qed
from *inf* **show** *inf'* $\sqsubseteq$ *inf*
proof (*rule is-inf-greatest*)
from *inf'* **show** *inf'* $\sqsubseteq x$ **..**
from *inf'* **show** *inf'* $\sqsubseteq y$ **..**
qed
qed
qed

theorem *is-sup-uniq*: *is-sup x y sup* $\Longrightarrow$ *is-sup x y sup'* $\Longrightarrow$ *sup* = *sup'*
proof −
  assume *sup*: *is-sup x y sup* **and** *sup'*: *is-sup x y sup'*
  have *dual sup* = *dual sup'*
  proof (*rule is-inf-uniq*)
    from *sup* **show** *is-inf* (*dual x*) (*dual y*) (*dual sup*) **..**
    from *sup'* **show** *is-inf* (*dual x*) (*dual y*) (*dual sup'*) **..**
  qed
  then show *sup* = *sup'* **..**
qed

theorem *is-Inf-uniq*: *is-Inf A inf* $\Longrightarrow$ *is-Inf A inf'* $\Longrightarrow$ *inf* = *inf'*
proof −
  assume *inf*: *is-Inf A inf*
  assume *inf'*: *is-Inf A inf'*
  show *?thesis*
  proof (*rule leq-antisym*)
    from *inf'* **show** *inf* $\sqsubseteq$ *inf'*
    proof (*rule is-Inf-greatest*)
      fix *x* **assume** *x* $\in$ *A*
      with *inf* **show** *inf* $\sqsubseteq$ *x* **..**
    qed
    from *inf* **show** *inf'* $\sqsubseteq$ *inf*
    proof (*rule is-Inf-greatest*)
      fix *x* **assume** *x* $\in$ *A*
      with *inf'* **show** *inf'* $\sqsubseteq$ *x* **..**
    qed
  qed
qed

theorem *is-Sup-uniq*: *is-Sup A sup* $\Longrightarrow$ *is-Sup A sup'* $\Longrightarrow$ *sup* = *sup'*
proof −
  assume *sup*: *is-Sup A sup* **and** *sup'*: *is-Sup A sup'*
  have *dual sup* = *dual sup'*
  proof (*rule is-Inf-uniq*)
    from *sup* **show** *is-Inf* (*dual ' A*) (*dual sup*) **..**
    from *sup'* **show** *is-Inf* (*dual ' A*) (*dual sup'*) **..**
  qed
  then show *sup* = *sup'* **..**
qed

2.4 Related elements

The binary bound of related elements is either one of the argument.

theorem *is-inf-related* [*elim?*]: *x* $\sqsubseteq$ *y* $\Longrightarrow$ *is-inf x y x*
proof −
  assume *x* $\sqsubseteq$ *y*
  show *?thesis*
  proof

```
    show x ⊑ x ..
    show x ⊑ y by fact
    fix z assume z ⊑ x and z ⊑ y show z ⊑ x by fact
  qed
qed

theorem is-sup-related [elim?]: x ⊑ y ⟹ is-sup x y y
proof −
  assume x ⊑ y
  show ?thesis
  proof
    show x ⊑ y by fact
    show y ⊑ y ..
    fix z assume x ⊑ z and y ⊑ z
    show y ⊑ z by fact
  qed
qed
```

2.5 General versus binary bounds

General bounds of two-element sets coincide with binary bounds.

```
theorem is-Inf-binary: is-Inf {x, y} inf = is-inf x y inf
proof −
  let ?A = {x, y}
  show ?thesis
  proof
    assume is-Inf: is-Inf ?A inf
    show is-inf x y inf
    proof
      have x ∈ ?A by simp
      with is-Inf show inf ⊑ x ..
      have y ∈ ?A by simp
      with is-Inf show inf ⊑ y ..
      fix z assume zx: z ⊑ x and zy: z ⊑ y
      from is-Inf show z ⊑ inf
      proof (rule is-Inf-greatest)
        fix a assume a ∈ ?A
        then have a = x ∨ a = y by blast
        then show z ⊑ a
        proof
          assume a = x
          with zx show ?thesis by simp
        next
          assume a = y
          with zy show ?thesis by simp
        qed
      qed
    qed
  next
```

```
    assume is-inf: is-inf x y inf
    show is-Inf {x, y} inf
    proof
      fix a assume a ∈ ?A
      then have a = x ∨ a = y by blast
      then show inf ⊑ a
      proof
        assume a = x
        also from is-inf have inf ⊑ x ..
        finally show ?thesis .
      next
        assume a = y
        also from is-inf have inf ⊑ y ..
        finally show ?thesis .
      qed
    next
      fix z assume z: ∀ a ∈ ?A. z ⊑ a
      from is-inf show z ⊑ inf
      proof (rule is-inf-greatest)
        from z show z ⊑ x by blast
        from z show z ⊑ y by blast
      qed
    qed
  qed
qed

theorem is-Sup-binary: is-Sup {x, y} sup = is-sup x y sup
proof −
  have is-Sup {x, y} sup = is-Inf (dual ' {x, y}) (dual sup)
    by (simp only: dual-Inf)
  also have dual ' {x, y} = {dual x, dual y}
    by simp
  also have is-Inf ... (dual sup) = is-inf (dual x) (dual y) (dual sup)
    by (rule is-Inf-binary)
  also have ... = is-sup x y sup
    by (simp only: dual-inf)
  finally show ?thesis .
qed
```

2.6 Connecting general bounds

Either kind of general bounds is sufficient to express the other. The least upper bound (supremum) is the same as the the greatest lower bound of the set of all upper bounds; the dual statements holds as well; the dual statement holds as well.

```
theorem Inf-Sup: is-Inf {b. ∀ a ∈ A. a ⊑ b} sup ⟹ is-Sup A sup
proof −
  let ?B = {b. ∀ a ∈ A. a ⊑ b}
```

```
  assume is-Inf: is-Inf ?B sup
  show is-Sup A sup
  proof
    fix x assume x: x ∈ A
    from is-Inf show x ⊑ sup
    proof (rule is-Inf-greatest)
      fix y assume y ∈ ?B
      then have ∀ a ∈ A. a ⊑ y ..
      from this x show x ⊑ y ..
    qed
  next
    fix z assume ∀ x ∈ A. x ⊑ z
    then have z ∈ ?B ..
    with is-Inf show sup ⊑ z ..
  qed
qed

theorem Sup-Inf: is-Sup {b. ∀ a ∈ A. b ⊑ a} inf ⟹ is-Inf A inf
proof −
  assume is-Sup {b. ∀ a ∈ A. b ⊑ a} inf
  then have is-Inf (dual ' {b. ∀ a ∈ A. dual a ⊑ dual b}) (dual inf)
    by (simp only: dual-Inf dual-leq)
  also have dual ' {b. ∀ a ∈ A. dual a ⊑ dual b} = {b'. ∀ a' ∈ dual ' A. a' ⊑ b'}
    by (auto iff: dual-ball dual-Collect simp add: image-Collect)
  finally have is-Inf ... (dual inf) .
  then have is-Sup (dual ' A) (dual inf)
    by (rule Inf-Sup)
  then show ?thesis ..
qed

end
```

3 Lattices

```
theory Lattice imports Bounds begin
```

3.1 Lattice operations

A *lattice* is a partial order with infimum and supremum of any two elements (thus any *finite* number of elements have bounds as well).

```
class lattice =
  assumes ex-inf: ∃ inf. is-inf x y inf
  assumes ex-sup: ∃ sup. is-sup x y sup
```

The ⊓ (meet) and ⊔ (join) operations select such infimum and supremum elements.

```
definition
  meet :: 'a::lattice ⇒ 'a ⇒ 'a  (infixl ‹⊓› 70) where
```

$x \sqcap y = (THE\ inf.\ is\text{-}inf\ x\ y\ inf)$
definition
$join :: 'a::lattice \Rightarrow 'a \Rightarrow 'a$ (**infixl** ‹⊔› 65) **where**
$x \sqcup y = (THE\ sup.\ is\text{-}sup\ x\ y\ sup)$

Due to unique existence of bounds, the lattice operations may be exhibited as follows.

lemma *meet-equality* [*elim?*]: $is\text{-}inf\ x\ y\ inf \Longrightarrow x \sqcap y = inf$
proof (*unfold meet-def*)
assume *is-inf x y inf*
then show $(THE\ inf.\ is\text{-}inf\ x\ y\ inf) = inf$
by (*rule the-equality*) (*rule is-inf-uniq* [*OF* - ‹*is-inf x y inf*›])
qed

lemma *meetI* [*intro?*]:
$inf \sqsubseteq x \Longrightarrow inf \sqsubseteq y \Longrightarrow (\bigwedge z.\ z \sqsubseteq x \Longrightarrow z \sqsubseteq y \Longrightarrow z \sqsubseteq inf) \Longrightarrow x \sqcap y = inf$
by (*rule meet-equality*, *rule is-infI*) *blast+*

lemma *join-equality* [*elim?*]: $is\text{-}sup\ x\ y\ sup \Longrightarrow x \sqcup y = sup$
proof (*unfold join-def*)
assume *is-sup x y sup*
then show $(THE\ sup.\ is\text{-}sup\ x\ y\ sup) = sup$
by (*rule the-equality*) (*rule is-sup-uniq* [*OF* - ‹*is-sup x y sup*›])
qed

lemma *joinI* [*intro?*]: $x \sqsubseteq sup \Longrightarrow y \sqsubseteq sup \Longrightarrow$
$(\bigwedge z.\ x \sqsubseteq z \Longrightarrow y \sqsubseteq z \Longrightarrow sup \sqsubseteq z) \Longrightarrow x \sqcup y = sup$
by (*rule join-equality*, *rule is-supI*) *blast+*

The $\sqcap$ and $\sqcup$ operations indeed determine bounds on a lattice structure.

lemma *is-inf-meet* [*intro?*]: $is\text{-}inf\ x\ y\ (x \sqcap y)$
proof (*unfold meet-def*)
from *ex-inf* **obtain** *inf* **where** *is-inf x y inf* **..**
then show $is\text{-}inf\ x\ y\ (THE\ inf.\ is\text{-}inf\ x\ y\ inf)$
by (*rule theI*) (*rule is-inf-uniq* [*OF* - ‹*is-inf x y inf*›])
qed

lemma *meet-greatest* [*intro?*]: $z \sqsubseteq x \Longrightarrow z \sqsubseteq y \Longrightarrow z \sqsubseteq x \sqcap y$
by (*rule is-inf-greatest*) (*rule is-inf-meet*)

lemma *meet-lower1* [*intro?*]: $x \sqcap y \sqsubseteq x$
by (*rule is-inf-lower*) (*rule is-inf-meet*)

lemma *meet-lower2* [*intro?*]: $x \sqcap y \sqsubseteq y$
by (*rule is-inf-lower*) (*rule is-inf-meet*)

lemma *is-sup-join* [*intro?*]: $is\text{-}sup\ x\ y\ (x \sqcup y)$

proof (*unfold join-def*)
  from *ex-sup* **obtain** *sup* **where** *is-sup x y sup* **..**
  then show *is-sup x y* (*THE sup. is-sup x y sup*)
    by (*rule theI*) (*rule is-sup-uniq* [*OF* - ‹*is-sup x y sup*›])
qed

lemma *join-least* [*intro?*]: $x \sqsubseteq z \Longrightarrow y \sqsubseteq z \Longrightarrow x \sqcup y \sqsubseteq z$
  by (*rule is-sup-least*) (*rule is-sup-join*)

lemma *join-upper1* [*intro?*]: $x \sqsubseteq x \sqcup y$
  by (*rule is-sup-upper*) (*rule is-sup-join*)

lemma *join-upper2* [*intro?*]: $y \sqsubseteq x \sqcup y$
  by (*rule is-sup-upper*) (*rule is-sup-join*)

3.2 Duality

The class of lattices is closed under formation of dual structures. This means that for any theorem of lattice theory, the dualized statement holds as well; this important fact simplifies many proofs of lattice theory.

instance *dual* :: (*lattice*) *lattice*
proof
  fix x' y' :: 'a::*lattice dual*
  show ∃ *inf'*. *is-inf x' y' inf'*
  proof −
    have ∃ *sup*. *is-sup* (*undual x'*) (*undual y'*) *sup* **by** (*rule ex-sup*)
    then have ∃ *sup*. *is-inf* (*dual* (*undual x'*)) (*dual* (*undual y'*)) (*dual sup*)
      by (*simp only*: *dual-inf*)
    then show *?thesis* **by** (*simp add*: *dual-ex* [*symmetric*])
  qed
  show ∃ *sup'*. *is-sup x' y' sup'*
  proof −
    have ∃ *inf*. *is-inf* (*undual x'*) (*undual y'*) *inf* **by** (*rule ex-inf*)
    then have ∃ *inf*. *is-sup* (*dual* (*undual x'*)) (*dual* (*undual y'*)) (*dual inf*)
      by (*simp only*: *dual-sup*)
    then show *?thesis* **by** (*simp add*: *dual-ex* [*symmetric*])
  qed
qed

Apparently, the $\sqcap$ and $\sqcup$ operations are dual to each other.

theorem *dual-meet* [*intro?*]: *dual* $(x \sqcap y)$ = *dual x* $\sqcup$ *dual y*
proof −
  from *is-inf-meet* **have** *is-sup* (*dual x*) (*dual y*) (*dual* $(x \sqcap y)$) **..**
  then have *dual x* $\sqcup$ *dual y* = *dual* $(x \sqcap y)$ **..**
  then show *?thesis* **..**
qed

theorem *dual-join* [*intro?*]: *dual* $(x \sqcup y)$ = *dual x* $\sqcap$ *dual y*

proof −
from *is-sup-join* **have** *is-inf* (*dual x*) (*dual y*) (*dual* (*x* ⊔ *y*)) **..**
then have *dual x* ⊓ *dual y* = *dual* (*x* ⊔ *y*) **..**
then show *?thesis* **..**
qed

3.3 Algebraic properties

The ⊓ and ⊔ operations have the following characteristic algebraic properties: associative (A), commutative (C), and absorptive (AB).

theorem *meet-assoc*: (*x* ⊓ *y*) ⊓ *z* = *x* ⊓ (*y* ⊓ *z*)
proof
show *x* ⊓ (*y* ⊓ *z*) ⊑ *x* ⊓ *y*
proof
show *x* ⊓ (*y* ⊓ *z*) ⊑ *x* **..**
show *x* ⊓ (*y* ⊓ *z*) ⊑ *y*
proof −
have *x* ⊓ (*y* ⊓ *z*) ⊑ *y* ⊓ *z* **..**
also have ... ⊑ *y* **..**
finally show *?thesis* **.**
qed
qed
show *x* ⊓ (*y* ⊓ *z*) ⊑ *z*
proof −
have *x* ⊓ (*y* ⊓ *z*) ⊑ *y* ⊓ *z* **..**
also have ... ⊑ *z* **..**
finally show *?thesis* **.**
qed
fix *w* **assume** *w* ⊑ *x* ⊓ *y* **and** *w* ⊑ *z*
show *w* ⊑ *x* ⊓ (*y* ⊓ *z*)
proof
show *w* ⊑ *x*
proof −
have *w* ⊑ *x* ⊓ *y* **by** *fact*
also have ... ⊑ *x* **..**
finally show *?thesis* **.**
qed
show *w* ⊑ *y* ⊓ *z*
proof
show *w* ⊑ *y*
proof −
have *w* ⊑ *x* ⊓ *y* **by** *fact*
also have ... ⊑ *y* **..**
finally show *?thesis* **.**
qed
show *w* ⊑ *z* **by** *fact*
qed
qed
qed

theorem *join-assoc*: $(x \sqcup y) \sqcup z = x \sqcup (y \sqcup z)$
proof −
  have *dual* $((x \sqcup y) \sqcup z) = (\mathit{dual}\ x \sqcap \mathit{dual}\ y) \sqcap \mathit{dual}\ z$
    by (*simp only*: *dual-join*)
  also have $\ldots = \mathit{dual}\ x \sqcap (\mathit{dual}\ y \sqcap \mathit{dual}\ z)$
    by (*rule meet-assoc*)
  also have $\ldots = \mathit{dual}\ (x \sqcup (y \sqcup z))$
    by (*simp only*: *dual-join*)
  finally show *?thesis* **..**
qed

theorem *meet-commute*: $x \sqcap y = y \sqcap x$
proof
  show $y \sqcap x \sqsubseteq x$ **..**
  show $y \sqcap x \sqsubseteq y$ **..**
  fix z **assume** $z \sqsubseteq y$ **and** $z \sqsubseteq x$
  then show $z \sqsubseteq y \sqcap x$ **..**
qed

theorem *join-commute*: $x \sqcup y = y \sqcup x$
proof −
  have *dual* $(x \sqcup y) = \mathit{dual}\ x \sqcap \mathit{dual}\ y$ **..**
  also have $\ldots = \mathit{dual}\ y \sqcap \mathit{dual}\ x$
    by (*rule meet-commute*)
  also have $\ldots = \mathit{dual}\ (y \sqcup x)$
    by (*simp only*: *dual-join*)
  finally show *?thesis* **..**
qed

theorem *meet-join-absorb*: $x \sqcap (x \sqcup y) = x$
proof
  show $x \sqsubseteq x$ **..**
  show $x \sqsubseteq x \sqcup y$ **..**
  fix z **assume** $z \sqsubseteq x$ **and** $z \sqsubseteq x \sqcup y$
  show $z \sqsubseteq x$ **by** *fact*
qed

theorem *join-meet-absorb*: $x \sqcup (x \sqcap y) = x$
proof −
  have $\mathit{dual}\ x \sqcap (\mathit{dual}\ x \sqcup \mathit{dual}\ y) = \mathit{dual}\ x$
    by (*rule meet-join-absorb*)
  then have *dual* $(x \sqcup (x \sqcap y)) = \mathit{dual}\ x$
    by (*simp only*: *dual-meet dual-join*)
  then show *?thesis* **..**
qed

Some further algebraic properties hold as well. The property idempotent (I) is a basic algebraic consequence of (AB).

theorem *meet-idem*: $x \sqcap x = x$
proof −
  have $x \sqcap (x \sqcup (x \sqcap x)) = x$ **by** (*rule meet-join-absorb*)
  also have $x \sqcup (x \sqcap x) = x$ **by** (*rule join-meet-absorb*)
  finally show *?thesis* **.**
qed

theorem *join-idem*: $x \sqcup x = x$
proof −
  have *dual* $x \sqcap$ *dual* $x =$ *dual* x
    by (*rule meet-idem*)
  then have *dual* $(x \sqcup x) =$ *dual* x
    by (*simp only*: *dual-join*)
  then show *?thesis* **..**
qed

Meet and join are trivial for related elements.

theorem *meet-related* [*elim?*]: $x \sqsubseteq y \Longrightarrow x \sqcap y = x$
proof
  assume $x \sqsubseteq y$
  show $x \sqsubseteq x$ **..**
  show $x \sqsubseteq y$ **by** *fact*
  fix z **assume** $z \sqsubseteq x$ **and** $z \sqsubseteq y$
  show $z \sqsubseteq x$ **by** *fact*
qed

theorem *join-related* [*elim?*]: $x \sqsubseteq y \Longrightarrow x \sqcup y = y$
proof −
  assume $x \sqsubseteq y$ **then have** *dual* $y \sqsubseteq$ *dual* x **..**
  then have *dual* $y \sqcap$ *dual* $x =$ *dual* y **by** (*rule meet-related*)
  also have *dual* $y \sqcap$ *dual* $x =$ *dual* $(y \sqcup x)$ **by** (*simp only*: *dual-join*)
  also have $y \sqcup x = x \sqcup y$ **by** (*rule join-commute*)
  finally show *?thesis* **..**
qed

3.4 Order versus algebraic structure

The $\sqcap$ and $\sqcup$ operations are connected with the underlying $\sqsubseteq$ relation in a canonical manner.

theorem *meet-connection*: $(x \sqsubseteq y) = (x \sqcap y = x)$
proof
  assume $x \sqsubseteq y$
  then have *is-inf* $x\ y\ x$ **..**
  then show $x \sqcap y = x$ **..**
next
  have $x \sqcap y \sqsubseteq y$ **..**
  also assume $x \sqcap y = x$
  finally show $x \sqsubseteq y$ **.**

qed

theorem *join-connection*: $(x \sqsubseteq y) = (x \sqcup y = y)$
proof
  assume $x \sqsubseteq y$
  then have *is-sup x y y* **..**
  then show $x \sqcup y = y$ **..**
next
  have $x \sqsubseteq x \sqcup y$ **..**
  also assume $x \sqcup y = y$
  finally show $x \sqsubseteq y$ **.**
qed

The most fundamental result of the meta-theory of lattices is as follows (we do not prove it here).

Given a structure with binary operations $\sqcap$ and $\sqcup$ such that (A), (C), and (AB) hold (cf. §3.3). This structure represents a lattice, if the relation $x \sqsubseteq y$ is defined as $x \sqcap y = x$ (alternatively as $x \sqcup y = y$). Furthermore, infimum and supremum with respect to this ordering coincide with the original $\sqcap$ and $\sqcup$ operations.

3.5 Example instances

3.5.1 Linear orders

Linear orders with *minimum* and *maximum* operations are a (degenerate) example of lattice structures.

definition
  minimum :: *'a*::*linear-order* ⇒ *'a* ⇒ *'a* **where**
  minimum x y = (*if x* ⊑ *y then x else y*)
definition
  maximum :: *'a*::*linear-order* ⇒ *'a* ⇒ *'a* **where**
  maximum x y = (*if x* ⊑ *y then y else x*)

lemma *is-inf-minimum*: *is-inf x y* (*minimum x y*)
proof
  let *?min* = *minimum x y*
  from *leq-linear* **show** *?min* ⊑ *x* **by** (*auto simp add*: *minimum-def*)
  from *leq-linear* **show** *?min* ⊑ *y* **by** (*auto simp add*: *minimum-def*)
  fix *z* **assume** *z* ⊑ *x* **and** *z* ⊑ *y*
  with *leq-linear* **show** *z* ⊑ *?min* **by** (*auto simp add*: *minimum-def*)
qed

lemma *is-sup-maximum*: *is-sup x y* (*maximum x y*)
proof
  let *?max* = *maximum x y*
  from *leq-linear* **show** *x* ⊑ *?max* **by** (*auto simp add*: *maximum-def*)

from *leq-linear* **show** $y \sqsubseteq ?max$ **by** (*auto simp add*: *maximum-def*)
fix z **assume** $x \sqsubseteq z$ **and** $y \sqsubseteq z$
with *leq-linear* **show** $?max \sqsubseteq z$ **by** (*auto simp add*: *maximum-def*)
qed

instance *linear-order* $\subseteq$ *lattice*
proof
fix $x\ y :: 'a$::*linear-order*
from *is-inf-minimum* **show** $\exists\, inf.$ *is-inf* $x\ y\ inf$ **..**
from *is-sup-maximum* **show** $\exists\, sup.$ *is-sup* $x\ y\ sup$ **..**
qed

The lattice operations on linear orders indeed coincide with *minimum* and *maximum*.

theorem *meet-mimimum*: $x \sqcap y = minimum\ x\ y$
by (*rule meet-equality*) (*rule is-inf-minimum*)

theorem *meet-maximum*: $x \sqcup y = maximum\ x\ y$
by (*rule join-equality*) (*rule is-sup-maximum*)

3.5.2 Binary products

The class of lattices is closed under direct binary products (cf. §1.3.2).

lemma *is-inf-prod*: *is-inf* $p\ q$ $(fst\ p \sqcap fst\ q,\ snd\ p \sqcap snd\ q)$
proof
show $(fst\ p \sqcap fst\ q,\ snd\ p \sqcap snd\ q) \sqsubseteq p$
proof −
have $fst\ p \sqcap fst\ q \sqsubseteq fst\ p$ **..**
moreover have $snd\ p \sqcap snd\ q \sqsubseteq snd\ p$ **..**
ultimately show *?thesis* **by** (*simp add*: *leq-prod-def*)
qed
show $(fst\ p \sqcap fst\ q,\ snd\ p \sqcap snd\ q) \sqsubseteq q$
proof −
have $fst\ p \sqcap fst\ q \sqsubseteq fst\ q$ **..**
moreover have $snd\ p \sqcap snd\ q \sqsubseteq snd\ q$ **..**
ultimately show *?thesis* **by** (*simp add*: *leq-prod-def*)
qed
fix r **assume** *rp*: $r \sqsubseteq p$ **and** *rq*: $r \sqsubseteq q$
show $r \sqsubseteq (fst\ p \sqcap fst\ q,\ snd\ p \sqcap snd\ q)$
proof −
have $fst\ r \sqsubseteq fst\ p \sqcap fst\ q$
proof
from *rp* **show** $fst\ r \sqsubseteq fst\ p$ **by** (*simp add*: *leq-prod-def*)
from *rq* **show** $fst\ r \sqsubseteq fst\ q$ **by** (*simp add*: *leq-prod-def*)
qed
moreover have $snd\ r \sqsubseteq snd\ p \sqcap snd\ q$
proof
from *rp* **show** $snd\ r \sqsubseteq snd\ p$ **by** (*simp add*: *leq-prod-def*)

```
    from rq show snd r ⊑ snd q by (simp add: leq-prod-def)
   qed
   ultimately show ?thesis by (simp add: leq-prod-def)
  qed
qed

lemma is-sup-prod: is-sup p q (fst p ⊔ fst q, snd p ⊔ snd q)
proof
  show p ⊑ (fst p ⊔ fst q, snd p ⊔ snd q)
  proof −
   have fst p ⊑ fst p ⊔ fst q ..
   moreover have snd p ⊑ snd p ⊔ snd q ..
   ultimately show ?thesis by (simp add: leq-prod-def)
  qed
  show q ⊑ (fst p ⊔ fst q, snd p ⊔ snd q)
  proof −
   have fst q ⊑ fst p ⊔ fst q ..
   moreover have snd q ⊑ snd p ⊔ snd q ..
   ultimately show ?thesis by (simp add: leq-prod-def)
  qed
  fix r assume pr: p ⊑ r and qr: q ⊑ r
  show (fst p ⊔ fst q, snd p ⊔ snd q) ⊑ r
  proof −
   have fst p ⊔ fst q ⊑ fst r
   proof
    from pr show fst p ⊑ fst r by (simp add: leq-prod-def)
    from qr show fst q ⊑ fst r by (simp add: leq-prod-def)
   qed
   moreover have snd p ⊔ snd q ⊑ snd r
   proof
    from pr show snd p ⊑ snd r by (simp add: leq-prod-def)
    from qr show snd q ⊑ snd r by (simp add: leq-prod-def)
   qed
   ultimately show ?thesis by (simp add: leq-prod-def)
  qed
qed

instance prod :: (lattice, lattice) lattice
proof
  fix p q :: 'a::lattice × 'b::lattice
  from is-inf-prod show ∃ inf. is-inf p q inf ..
  from is-sup-prod show ∃ sup. is-sup p q sup ..
qed
```

The lattice operations on a binary product structure indeed coincide with the products of the original ones.

```
theorem meet-prod: p ⊓ q = (fst p ⊓ fst q, snd p ⊓ snd q)
  by (rule meet-equality) (rule is-inf-prod)
```

theorem *join-prod*: $p \sqcup q = (fst\ p \sqcup fst\ q,\ snd\ p \sqcup snd\ q)$
by (*rule join-equality*) (*rule is-sup-prod*)

3.5.3 General products

The class of lattices is closed under general products (function spaces) as well (cf. §1.3.3).

lemma *is-inf-fun*: *is-inf f g* $(\lambda x.\ f\ x \sqcap g\ x)$
proof
show $(\lambda x.\ f\ x \sqcap g\ x) \sqsubseteq f$
proof
fix x **show** $f\ x \sqcap g\ x \sqsubseteq f\ x$ **..**
qed
show $(\lambda x.\ f\ x \sqcap g\ x) \sqsubseteq g$
proof
fix x **show** $f\ x \sqcap g\ x \sqsubseteq g\ x$ **..**
qed
fix h **assume** *hf*: $h \sqsubseteq f$ **and** *hg*: $h \sqsubseteq g$
show $h \sqsubseteq (\lambda x.\ f\ x \sqcap g\ x)$
proof
fix x
show $h\ x \sqsubseteq f\ x \sqcap g\ x$
proof
from *hf* **show** $h\ x \sqsubseteq f\ x$ **..**
from *hg* **show** $h\ x \sqsubseteq g\ x$ **..**
qed
qed
qed

lemma *is-sup-fun*: *is-sup f g* $(\lambda x.\ f\ x \sqcup g\ x)$
proof
show $f \sqsubseteq (\lambda x.\ f\ x \sqcup g\ x)$
proof
fix x **show** $f\ x \sqsubseteq f\ x \sqcup g\ x$ **..**
qed
show $g \sqsubseteq (\lambda x.\ f\ x \sqcup g\ x)$
proof
fix x **show** $g\ x \sqsubseteq f\ x \sqcup g\ x$ **..**
qed
fix h **assume** *fh*: $f \sqsubseteq h$ **and** *gh*: $g \sqsubseteq h$
show $(\lambda x.\ f\ x \sqcup g\ x) \sqsubseteq h$
proof
fix x
show $f\ x \sqcup g\ x \sqsubseteq h\ x$
proof
from *fh* **show** $f\ x \sqsubseteq h\ x$ **..**
from *gh* **show** $g\ x \sqsubseteq h\ x$ **..**
qed
qed

qed

instance *fun* :: (*type*, *lattice*) *lattice*
proof
  fix *f g* :: *'a* ⇒ *'b*::*lattice*
  show ∃ *inf*. *is-inf f g inf* **by** *rule* (*rule is-inf-fun*)
  show ∃ *sup*. *is-sup f g sup* **by** *rule* (*rule is-sup-fun*)
qed

The lattice operations on a general product structure (function space) indeed emerge by point-wise lifting of the original ones.

theorem *meet-fun*: *f* ⊓ *g* = (λ*x*. *f x* ⊓ *g x*)
  by (*rule meet-equality*) (*rule is-inf-fun*)

theorem *join-fun*: *f* ⊔ *g* = (λ*x*. *f x* ⊔ *g x*)
  by (*rule join-equality*) (*rule is-sup-fun*)

3.6 Monotonicity and semi-morphisms

The lattice operations are monotone in both argument positions. In fact, monotonicity of the second position is trivial due to commutativity.

theorem *meet-mono*: *x* ⊑ *z* ⟹ *y* ⊑ *w* ⟹ *x* ⊓ *y* ⊑ *z* ⊓ *w*
proof −
  {
    fix *a b c* :: *'a*::*lattice*
    assume *a* ⊑ *c* **have** *a* ⊓ *b* ⊑ *c* ⊓ *b*
    proof
      have *a* ⊓ *b* ⊑ *a* **..**
      also have ... ⊑ *c* **by** *fact*
      finally show *a* ⊓ *b* ⊑ *c* **.**
      show *a* ⊓ *b* ⊑ *b* **..**
    qed
  } note *this* [*elim?*]
  assume *x* ⊑ *z* **then have** *x* ⊓ *y* ⊑ *z* ⊓ *y* **..**
  also have ... = *y* ⊓ *z* **by** (*rule meet-commute*)
  also assume *y* ⊑ *w* **then have** *y* ⊓ *z* ⊑ *w* ⊓ *z* **..**
  also have ... = *z* ⊓ *w* **by** (*rule meet-commute*)
  finally show *?thesis* **.**
qed

theorem *join-mono*: *x* ⊑ *z* ⟹ *y* ⊑ *w* ⟹ *x* ⊔ *y* ⊑ *z* ⊔ *w*
proof −
  assume *x* ⊑ *z* **then have** *dual z* ⊑ *dual x* **..**
  moreover assume *y* ⊑ *w* **then have** *dual w* ⊑ *dual y* **..**
  ultimately have *dual z* ⊓ *dual w* ⊑ *dual x* ⊓ *dual y*
    by (*rule meet-mono*)
  then have *dual* (*z* ⊔ *w*) ⊑ *dual* (*x* ⊔ *y*)
    by (*simp only*: *dual-join*)

then show *?thesis* **..**
qed

A semi-morphisms is a function f that preserves the lattice operations in the following manner: $f\ (x \sqcap y) \sqsubseteq f\ x \sqcap f\ y$ and $f\ x \sqcup f\ y \sqsubseteq f\ (x \sqcup y)$, respectively. Any of these properties is equivalent with monotonicity.

theorem *meet-semimorph*:
$(\bigwedge x\ y.\ f\ (x \sqcap y) \sqsubseteq f\ x \sqcap f\ y) \equiv (\bigwedge x\ y.\ x \sqsubseteq y \Longrightarrow f\ x \sqsubseteq f\ y)$
proof
assume *morph*: $\bigwedge x\ y.\ f\ (x \sqcap y) \sqsubseteq f\ x \sqcap f\ y$
fix $x\ y :: 'a{::}lattice$
assume $x \sqsubseteq y$
then have $x \sqcap y = x$ **..**
then have $x = x \sqcap y$ **..**
also have $f\ \ldots \sqsubseteq f\ x \sqcap f\ y$ **by** (*rule morph*)
also have $\ldots \sqsubseteq f\ y$ **..**
finally show $f\ x \sqsubseteq f\ y$ **.**
next
assume *mono*: $\bigwedge x\ y.\ x \sqsubseteq y \Longrightarrow f\ x \sqsubseteq f\ y$
show $\bigwedge x\ y.\ f\ (x \sqcap y) \sqsubseteq f\ x \sqcap f\ y$
proof −
fix $x\ y$
show $f\ (x \sqcap y) \sqsubseteq f\ x \sqcap f\ y$
proof
have $x \sqcap y \sqsubseteq x$ **.. then show** $f\ (x \sqcap y) \sqsubseteq f\ x$ **by** (*rule mono*)
have $x \sqcap y \sqsubseteq y$ **.. then show** $f\ (x \sqcap y) \sqsubseteq f\ y$ **by** (*rule mono*)
qed
qed
qed

lemma *join-semimorph*:
$(\bigwedge x\ y.\ f\ x \sqcup f\ y \sqsubseteq f\ (x \sqcup y)) \equiv (\bigwedge x\ y.\ x \sqsubseteq y \Longrightarrow f\ x \sqsubseteq f\ y)$
proof
assume *morph*: $\bigwedge x\ y.\ f\ x \sqcup f\ y \sqsubseteq f\ (x \sqcup y)$
fix $x\ y :: 'a{::}lattice$
assume $x \sqsubseteq y$ **then have** $x \sqcup y = y$ **..**
have $f\ x \sqsubseteq f\ x \sqcup f\ y$ **..**
also have $\ldots \sqsubseteq f\ (x \sqcup y)$ **by** (*rule morph*)
also from $\langle x \sqsubseteq y \rangle$ **have** $x \sqcup y = y$ **..**
finally show $f\ x \sqsubseteq f\ y$ **.**
next
assume *mono*: $\bigwedge x\ y.\ x \sqsubseteq y \Longrightarrow f\ x \sqsubseteq f\ y$
show $\bigwedge x\ y.\ f\ x \sqcup f\ y \sqsubseteq f\ (x \sqcup y)$
proof −
fix $x\ y$
show $f\ x \sqcup f\ y \sqsubseteq f\ (x \sqcup y)$
proof
have $x \sqsubseteq x \sqcup y$ **.. then show** $f\ x \sqsubseteq f\ (x \sqcup y)$ **by** (*rule mono*)
have $y \sqsubseteq x \sqcup y$ **.. then show** $f\ y \sqsubseteq f\ (x \sqcup y)$ **by** (*rule mono*)

qed
qed
qed

end

4 Complete lattices

theory *CompleteLattice* **imports** *Lattice* **begin**

4.1 Complete lattice operations

A *complete lattice* is a partial order with general (infinitary) infimum of any set of elements. General supremum exists as well, as a consequence of the connection of infinitary bounds (see §2.6).

class *complete-lattice* =
assumes *ex-Inf*: $\exists\, inf.$ *is-Inf A inf*

theorem *ex-Sup*: $\exists\, sup{::}'a{::}$*complete-lattice*. *is-Sup A sup*
proof −
from *ex-Inf* **obtain** *sup* **where** *is-Inf* $\{b.\ \forall\, a{\in}A.\ a \sqsubseteq b\}$ *sup* **by** *blast*
then have *is-Sup A sup* **by** (*rule Inf-Sup*)
then show *?thesis* **..**
qed

The general $\bigsqcap$ (meet) and $\bigsqcup$ (join) operations select such infimum and supremum elements.

definition
Meet :: $'a{::}$*complete-lattice set* ⇒ $'a$ (‹$\bigsqcap$-› [*90*] *90*) **where**
$\bigsqcap A = (THE\ inf.\ is\text{-}Inf\ A\ inf)$
definition
Join :: $'a{::}$*complete-lattice set* ⇒ $'a$ (‹$\bigsqcup$-› [*90*] *90*) **where**
$\bigsqcup A = (THE\ sup.\ is\text{-}Sup\ A\ sup)$

Due to unique existence of bounds, the complete lattice operations may be exhibited as follows.

lemma *Meet-equality* [*elim?*]: *is-Inf A inf* ⟹ $\bigsqcap A = inf$
proof (*unfold Meet-def*)
assume *is-Inf A inf*
then show (*THE inf*. *is-Inf A inf*) = *inf*
by (*rule the-equality*) (*rule is-Inf-uniq* [*OF* - ‹*is-Inf A inf*›])
qed

lemma *MeetI* [*intro?*]:
$(\bigwedge a.\ a \in A \Longrightarrow inf \sqsubseteq a) \Longrightarrow$
$(\bigwedge b.\ \forall\, a \in A.\ b \sqsubseteq a \Longrightarrow b \sqsubseteq inf) \Longrightarrow$
$\bigsqcap A = inf$

by (*rule Meet-equality*, *rule is-InfI*) *blast+*

lemma *Join-equality* [*elim?*]: *is-Sup A sup* $\Longrightarrow \bigsqcup A = sup$
proof (*unfold Join-def*)
assume *is-Sup A sup*
then show (*THE sup. is-Sup A sup*) = *sup*
by (*rule the-equality*) (*rule is-Sup-uniq* [*OF* - ‹*is-Sup A sup*›])
qed

lemma *JoinI* [*intro?*]:
$(\bigwedge a.\ a \in A \Longrightarrow a \sqsubseteq sup) \Longrightarrow$
$(\bigwedge b.\ \forall a \in A.\ a \sqsubseteq b \Longrightarrow sup \sqsubseteq b) \Longrightarrow$
$\bigsqcup A = sup$
by (*rule Join-equality*, *rule is-SupI*) *blast+*

The $\bigsqcap$ and $\bigsqcup$ operations indeed determine bounds on a complete lattice structure.

lemma *is-Inf-Meet* [*intro?*]: *is-Inf A* ($\bigsqcap A$)
proof (*unfold Meet-def*)
from *ex-Inf* **obtain** *inf* **where** *is-Inf A inf* **..**
then show *is-Inf A* (*THE inf*. *is-Inf A inf*)
by (*rule theI*) (*rule is-Inf-uniq* [*OF* - ‹*is-Inf A inf*›])
qed

lemma *Meet-greatest* [*intro?*]: $(\bigwedge a.\ a \in A \Longrightarrow x \sqsubseteq a) \Longrightarrow x \sqsubseteq \bigsqcap A$
by (*rule is-Inf-greatest*, *rule is-Inf-Meet*) *blast*

lemma *Meet-lower* [*intro?*]: $a \in A \Longrightarrow \bigsqcap A \sqsubseteq a$
by (*rule is-Inf-lower*) (*rule is-Inf-Meet*)

lemma *is-Sup-Join* [*intro?*]: *is-Sup A* ($\bigsqcup A$)
proof (*unfold Join-def*)
from *ex-Sup* **obtain** *sup* **where** *is-Sup A sup* **..**
then show *is-Sup A* (*THE sup. is-Sup A sup*)
by (*rule theI*) (*rule is-Sup-uniq* [*OF* - ‹*is-Sup A sup*›])
qed

lemma *Join-least* [*intro?*]: $(\bigwedge a.\ a \in A \Longrightarrow a \sqsubseteq x) \Longrightarrow \bigsqcup A \sqsubseteq x$
by (*rule is-Sup-least*, *rule is-Sup-Join*) *blast*
lemma *Join-lower* [*intro?*]: $a \in A \Longrightarrow a \sqsubseteq \bigsqcup A$
by (*rule is-Sup-upper*) (*rule is-Sup-Join*)

4.2 The Knaster-Tarski Theorem

The Knaster-Tarski Theorem (in its simplest formulation) states that any monotone function on a complete lattice has a least fixed-point (see [2, pages 93–94] for example). This is a consequence of the basic boundary properties

of the complete lattice operations.

theorem *Knaster-Tarski*:
assumes *mono*: $\bigwedge x\ y.\ x \sqsubseteq y \Longrightarrow f\ x \sqsubseteq f\ y$
obtains $a :: 'a$::*complete-lattice* **where**
$f\ a = a$ **and** $\bigwedge a'.\ f\ a' = a' \Longrightarrow a \sqsubseteq a'$
proof
let *?H* $= \{u.\ f\ u \sqsubseteq u\}$
let *?a* $= \bigsqcap$ *?H*
show *f ?a* $=$ *?a*
proof −
have *ge*: *f ?a* $\sqsubseteq$ *?a*
proof
fix *x* **assume** *x*: $x \in$ *?H*
then have *?a* $\sqsubseteq x$ **..**
then have *f ?a* $\sqsubseteq f\ x$ **by** (*rule mono*)
also from *x* **have** $\ldots \sqsubseteq x$ **..**
finally show *f ?a* $\sqsubseteq x$ **.**
qed
also have *?a* $\sqsubseteq$ *f ?a*
proof
from *ge* **have** *f* (*f ?a*) $\sqsubseteq$ *f ?a* **by** (*rule mono*)
then show *f ?a* $\in$ *?H* **..**
qed
finally show *?thesis* **.**
qed

fix a'
assume $f\ a' = a'$
then have $f\ a' \sqsubseteq a'$ **by** (*simp only*: *leq-refl*)
then have $a' \in$ *?H* **..**
then show *?a* $\sqsubseteq a'$ **..**
qed

theorem *Knaster-Tarski-dual*:
assumes *mono*: $\bigwedge x\ y.\ x \sqsubseteq y \Longrightarrow f\ x \sqsubseteq f\ y$
obtains $a :: 'a$::*complete-lattice* **where**
$f\ a = a$ **and** $\bigwedge a'.\ f\ a' = a' \Longrightarrow a' \sqsubseteq a$
proof
let *?H* $= \{u.\ u \sqsubseteq f\ u\}$
let *?a* $= \bigsqcup$ *?H*
show *f ?a* $=$ *?a*
proof −
have *le*: *?a* $\sqsubseteq$ *f ?a*
proof
fix *x* **assume** *x*: $x \in$ *?H*
then have $x \sqsubseteq f\ x$ **..**
also from *x* **have** $x \sqsubseteq$ *?a* **..**
then have $f\ x \sqsubseteq$ *f ?a* **by** (*rule mono*)
finally show $x \sqsubseteq$ *f ?a* **.**

```
    qed
    have f ?a ⊑ ?a
    proof
      from le have f ?a ⊑ f (f ?a) by (rule mono)
      then show f ?a ∈ ?H ..
    qed
    from this and le show ?thesis by (rule leq-antisym)
  qed

  fix a'
  assume f a' = a'
  then have a' ⊑ f a' by (simp only: leq-refl)
  then have a' ∈ ?H ..
  then show a' ⊑ ?a ..
qed
```

4.3 Bottom and top elements

With general bounds available, complete lattices also have least and greatest elements.

```
definition
  bottom :: 'a::complete-lattice  (‹⊥›) where
  ⊥ = ⨅ UNIV

definition
  top :: 'a::complete-lattice  (‹⊤›) where
  ⊤ = ⨆ UNIV

lemma bottom-least [intro?]: ⊥ ⊑ x
proof (unfold bottom-def)
  have x ∈ UNIV ..
  then show ⨅ UNIV ⊑ x ..
qed

lemma bottomI [intro?]: (⋀a. x ⊑ a) ⟹ ⊥ = x
proof (unfold bottom-def)
  assume ⋀a. x ⊑ a
  show ⨅ UNIV = x
  proof
    fix a show x ⊑ a by fact
  next
    fix b :: 'a::complete-lattice
    assume b: ∀ a ∈ UNIV. b ⊑ a
    have x ∈ UNIV ..
    with b show b ⊑ x ..
  qed
qed

lemma top-greatest [intro?]: x ⊑ ⊤
```

proof (*unfold top-def*)
  have $x \in \mathit{UNIV}$ **..**
  then show $x \sqsubseteq \bigsqcup \mathit{UNIV}$ **..**
qed

lemma *topI* [*intro?*]: $(\bigwedge a.\ a \sqsubseteq x) \Longrightarrow \top = x$
proof (*unfold top-def*)
  assume $\bigwedge a.\ a \sqsubseteq x$
  show $\bigsqcup \mathit{UNIV} = x$
  proof
    fix a **show** $a \sqsubseteq x$ **by** *fact*
  next
    fix $b :: 'a$::*complete-lattice*
    assume b: $\forall\, a \in \mathit{UNIV}.\ a \sqsubseteq b$
    have $x \in \mathit{UNIV}$ **..**
    with b **show** $x \sqsubseteq b$ **..**
  qed
qed

4.4 Duality

The class of complete lattices is closed under formation of dual structures.

instance *dual* :: (*complete-lattice*) *complete-lattice*
proof
  fix $A' :: 'a$::*complete-lattice dual set*
  show $\exists\, \mathit{inf}'.\ \mathit{is\text{-}Inf}\ A'\ \mathit{inf}'$
  proof −
    have $\exists\, \mathit{sup}.\ \mathit{is\text{-}Sup}\ (\mathit{undual}\ `\ A')\ \mathit{sup}$ **by** (*rule ex-Sup*)
   then have $\exists\, \mathit{sup}.\ \mathit{is\text{-}Inf}\ (\mathit{dual}\ `\ \mathit{undual}\ `\ A')\ (\mathit{dual}\ \mathit{sup})$ **by** (*simp only*: *dual-Inf*)
    then show *?thesis* **by** (*simp add*: *dual-ex* [*symmetric*] *image-comp*)
  qed
qed

Apparently, the $\bigsqcap$ and $\bigsqcup$ operations are dual to each other.

theorem *dual-Meet* [*intro?*]: $\mathit{dual}\ (\bigsqcap A) = \bigsqcup (\mathit{dual}\ `\ A)$
proof −
  from *is-Inf-Meet* **have** $\mathit{is\text{-}Sup}\ (\mathit{dual}\ `\ A)\ (\mathit{dual}\ (\bigsqcap A))$ **..**
  then have $\bigsqcup (\mathit{dual}\ `\ A) = \mathit{dual}\ (\bigsqcap A)$ **..**
  then show *?thesis* **..**
qed

theorem *dual-Join* [*intro?*]: $\mathit{dual}\ (\bigsqcup A) = \bigsqcap (\mathit{dual}\ `\ A)$
proof −
  from *is-Sup-Join* **have** $\mathit{is\text{-}Inf}\ (\mathit{dual}\ `\ A)\ (\mathit{dual}\ (\bigsqcup A))$ **..**
  then have $\bigsqcap (\mathit{dual}\ `\ A) = \mathit{dual}\ (\bigsqcup A)$ **..**
  then show *?thesis* **..**
qed

Likewise are $\bot$ and $\top$ duals of each other.

theorem *dual-bottom* [*intro?*]: *dual* $\bot = \top$
proof −
  have $\top =$ *dual* $\bot$
  proof
    fix a' **have** $\bot \sqsubseteq$ *undual* a' **..**
    then have *dual* (*undual* a') $\sqsubseteq$ *dual* $\bot$ **..**
    then show $a' \sqsubseteq$ *dual* $\bot$ **by** *simp*
  qed
  then show *?thesis* **..**
qed

theorem *dual-top* [*intro?*]: *dual* $\top = \bot$
proof −
  have $\bot =$ *dual* $\top$
  proof
    fix a' **have** *undual* $a' \sqsubseteq \top$ **..**
    then have *dual* $\top \sqsubseteq$ *dual* (*undual* a') **..**
    then show *dual* $\top \sqsubseteq a'$ **by** *simp*
  qed
  then show *?thesis* **..**
qed

4.5 Complete lattices are lattices

Complete lattices (with general bounds available) are indeed plain lattices as well. This holds due to the connection of general versus binary bounds that has been formally established in §2.5.

lemma *is-inf-binary*: *is-inf* x y ($\bigsqcap \{x, y\}$)
proof −
  have *is-Inf* $\{x, y\}$ ($\bigsqcap \{x, y\}$) **..**
  then show *?thesis* **by** (*simp only*: *is-Inf-binary*)
qed

lemma *is-sup-binary*: *is-sup* x y ($\bigsqcup \{x, y\}$)
proof −
  have *is-Sup* $\{x, y\}$ ($\bigsqcup \{x, y\}$) **..**
  then show *?thesis* **by** (*simp only*: *is-Sup-binary*)
qed

instance *complete-lattice* $\subseteq$ *lattice*
proof
  fix x y :: $'a$::*complete-lattice*
  from *is-inf-binary* **show** $\exists$ *inf*. *is-inf* x y *inf* **..**
  from *is-sup-binary* **show** $\exists$ *sup*. *is-sup* x y *sup* **..**
qed

theorem *meet-binary*: $x \sqcap y = \bigsqcap \{x, y\}$
  by (*rule meet-equality*) (*rule is-inf-binary*)

theorem *join-binary*: $x \sqcup y = \bigsqcup \{x, y\}$
by (*rule join-equality*) (*rule is-sup-binary*)

4.6 Complete lattices and set-theory operations

The complete lattice operations are (anti) monotone wrt. set inclusion.

theorem *Meet-subset-antimono*: $A \subseteq B \Longrightarrow \bigsqcap B \sqsubseteq \bigsqcap A$
proof (*rule Meet-greatest*)
fix a **assume** $a \in A$
also assume $A \subseteq B$
finally have $a \in B$ **.**
then show $\bigsqcap B \sqsubseteq a$ **..**
qed

theorem *Join-subset-mono*: $A \subseteq B \Longrightarrow \bigsqcup A \sqsubseteq \bigsqcup B$
proof −
assume $A \subseteq B$
then have *dual* ' $A \subseteq$ *dual* ' B **by** *blast*
then have $\bigsqcap$(*dual* ' B) $\sqsubseteq \bigsqcap$(*dual* ' A) **by** (*rule Meet-subset-antimono*)
then have *dual* ($\bigsqcup B$) $\sqsubseteq$ *dual* ($\bigsqcup A$) **by** (*simp only*: *dual-Join*)
then show *?thesis* **by** (*simp only*: *dual-leq*)
qed

Bounds over unions of sets may be obtained separately.

theorem *Meet-Un*: $\bigsqcap(A \cup B) = \bigsqcap A \sqcap \bigsqcap B$
proof
fix a **assume** $a \in A \cup B$
then show $\bigsqcap A \sqcap \bigsqcap B \sqsubseteq a$
proof
assume *a*: $a \in A$
have $\bigsqcap A \sqcap \bigsqcap B \sqsubseteq \bigsqcap A$ **..**
also from a **have** $\ldots \sqsubseteq a$ **..**
finally show *?thesis* **.**
next
assume *a*: $a \in B$
have $\bigsqcap A \sqcap \bigsqcap B \sqsubseteq \bigsqcap B$ **..**
also from a **have** $\ldots \sqsubseteq a$ **..**
finally show *?thesis* **.**
qed
next
fix b **assume** b: $\forall a \in A \cup B.\ b \sqsubseteq a$
show $b \sqsubseteq \bigsqcap A \sqcap \bigsqcap B$
proof
show $b \sqsubseteq \bigsqcap A$
proof
fix a **assume** $a \in A$
then have $a \in A \cup B$ **..**

```
      with b show b ⊑ a ..
    qed
    show b ⊑ ⨅ B
    proof
      fix a assume a ∈ B
      then have a ∈  A ∪ B ..
      with b show b ⊑ a ..
    qed
  qed
qed

theorem Join-Un: ⨆(A ∪ B) = ⨆ A ⊔ ⨆ B
proof −
  have dual (⨆(A ∪ B)) = ⨅(dual ‘ A ∪ dual ‘ B)
    by (simp only: dual-Join image-Un)
  also have ... = ⨅(dual ‘ A) ⊓ ⨅(dual ‘ B)
    by (rule Meet-Un)
  also have ... = dual (⨆ A ⊔ ⨆ B)
    by (simp only: dual-join dual-Join)
  finally show ?thesis ..
qed
```

Bounds over singleton sets are trivial.

```
theorem Meet-singleton: ⨅{x} = x
proof
  fix a assume a ∈ {x}
  then have a = x by simp
  then show x ⊑ a by (simp only: leq-refl)
next
  fix b assume ∀ a ∈ {x}. b ⊑ a
  then show b ⊑ x by simp
qed

theorem Join-singleton: ⨆{x} = x
proof −
  have dual (⨆{x}) = ⨅{dual x} by (simp add: dual-Join)
  also have ... = dual x by (rule Meet-singleton)
  finally show ?thesis ..
qed
```

Bounds over the empty and universal set correspond to each other.

```
theorem Meet-empty: ⨅{} = ⨆ UNIV
proof
  fix a :: 'a::complete-lattice
  assume a ∈ {}
  then have False by simp
  then show ⨆ UNIV ⊑ a ..
next
  fix b :: 'a::complete-lattice
```

```
  have b ∈ UNIV ..
  then show b ⊑ ⨆ UNIV ..
qed

theorem Join-empty: ⨆ {} = ⨅ UNIV
proof −
  have dual (⨆ {}) = ⨅ {} by (simp add: dual-Join)
  also have ... = ⨆ UNIV by (rule Meet-empty)
  also have ... = dual (⨅ UNIV) by (simp add: dual-Meet)
  finally show ?thesis ..
qed

end
```

References

[1] G. Bauer and M. Wenzel. Computer-assisted mathematics at work — the Hahn-Banach theorem in Isabelle/Isar. In T. Coquand, P. Dybjer, B. Nordström, and J. Smith, editors, *Types for Proofs and Programs: TYPES'99*, LNCS, 2000.

[2] B. A. Davey and H. A. Priestley. *Introduction to Lattices and Order*. Cambridge University Press, 1990.

[3] M. Wenzel. Isar — a generic interpretative approach to readable formal proof documents. In Y. Bertot, G. Dowek, A. Hirschowitz, C. Paulin, and L. Thery, editors, *Theorem Proving in Higher Order Logics: TPHOLs '99*, volume 1690 of *LNCS*, 1999.

[4] M. Wenzel. *The Isabelle/Isar Reference Manual*, 2000. https://isabelle.in.tum.de/doc/isar-ref.pdf.

[5] M. Wenzel. *Using Axiomatic Type Classes in Isabelle*, 2000. https://isabelle.in.tum.de/doc/axclass.pdf.